

Chem 4A Scholars Worksheet 2

Ionic Bonding and Electronegativity

Equations and useful constants:


$$F = -k \frac{q_1 q_2}{r^2} = -\frac{q_1 q_2}{4\pi\epsilon_0 r^2} \quad U = k \frac{q_1 q_2}{r} = \frac{q_1 q_2}{4\pi\epsilon_0 r} \quad e = 1.609 \times 10^{-19} \text{ C}$$

$$k = 8.988 \times 10^9 \text{ C}^{-2} \text{ N m}^2 \quad \epsilon_0 = 8.854 \times 10^{-12} \text{ C}^2 \text{ N}^{-1} \text{ m}^{-2}$$

Ionic Bonding:

Questions:

1. Warm up: Calculate the energy of a pair of Ca^{2+} and O^{2-} separated by 10.0 \AA .
2. If the above two ions are brought together to 5.00 \AA , will energy be released or absorbed. Find the energy.
3. Infinitely long NaCl chain. Imagine there are an infinitely long chain of alternating Na^+ and Cl^- , each separated by distance r , (i) write an expression for the potential energy felt by a Na^+ atom in the chain in r and necessary constant. Try use summation symbol to have a clear representation. (Extra Credit: If you have learned Taylor expansion in AP Calculus, try to write an analytical expression for the series) (ii) if $r=2.80 \text{ \AA}$, calculate the potential in J.



Electronegativity:

4. Rank the following elements by their electronegativity from lowest to highest: Na, Br, Fe, C, O, F, K, Cs.
5. Rank the the above elements in Ionization energy.
6. Is EA positive or negative?