


Chem 4A Scholars Worksheet 5

Blackbody Radiation, Bohr's Model, Spectroscopy, DeBroglie Relations

Useful equations:


$$\lambda_{max}T = b \quad \lambda = \frac{h}{p} = \frac{h}{mv} \quad v = -R_{\infty} \frac{Z^2}{n^2} c = 3.00 \times 10^8 \text{ m s}^{-1}$$

$$h = 6.63 \times 10^{-34} \text{ J s} \quad R_{\infty} = 10973731.6 \text{ m}^{-1} \quad 1 \text{ eV} = 1.602 \times 10^{-19} \text{ J} \quad m_e = 9.10 \times 10^{-31} \text{ kg}$$




Questions:

- Rank the following waves from lowest energy to highest energy. (Write 1 to 7 on top)
 radio wave, microwave, X-ray, gamma ray, infrared, ultraviolet, visible
- Fill in temperatures in the following blackbody radiation graph: 5500K, 4500K, 4000K, 3500K



- Blue Giants are a type of stars that glow blueish-white light, their maximum emission peak sit in the UV region. From the top graph, calculate the surface temperature of a star whose maximum emission is at 300nm.

4. Solve the following problem according to the Bohr's model:



A hydrogen atom was initially at some unknown energy level n_1 . The hydrogen atom absorbed a photon with wavelength of 434 nm , and went to a higher energy state n_3 . It then went through 2 transitions and emits two photons. The second photon was red. Find the wavelength of the first photon and the second photon was red. Find the energy levels n_1 , n_2 , n_3 , and the wavelength of the photons.

5. Would you expect a photon with wavenumber of 30000 cm^{-1} to be absorbed by a hydrogen atom at ground state?
6. Calculate the deBroglie wavelength of an electron whose speed is 500000 m s^{-1} .
7. Which requires more energy? Breaking CaCl_2 or Breaking CaO to ions?